

Orthogonal and Symplectic Matrix Integrals and Coupled KP Hierarchy

Saburo Kakei*

*Department of Mathematical Sciences, University of Tokyo,
Komaba 3-8-1, Meguro-ku, Tokyo 153-8902, Japan*

Abstract

Orthogonal and symplectic matrix integrals are investigated. It is shown that the matrix integrals can be considered as a τ -function of the coupled KP hierarchy, whose solution can be expressed in terms of pfaffians.

Over the past decade, the intimate relationships between matrix integrals and nonlinear integrable systems have been clarified, particularly in the context of string theory. In such cases, nonperturbative properties of physical quantities can be evaluated by the use of the integrable structures of the models. (For review, see refs. 1-4.)

Here, we consider a matrix integral over an ensemble of Hermitian matrices

$$Z_N^{(\beta)}\{t\} = \frac{1}{N!} \int_{-\infty}^{\infty} \cdots \int_{-\infty}^{\infty} \prod_{1 \leq j < k \leq N} |x_j - x_k|^\beta \exp \left[\sum_{i=1}^N \eta(x_i, t) \right] dx_1 \cdots dx_N, \quad (1)$$

where $\beta = 1, 2$ or 4 , for which the ensemble is, respectively, orthogonal, unitary or symplectic and $\eta(x, t) = \sum_{m=1}^{\infty} x^m t_m$. An interesting observation is that, when $\beta = 2$, the quantity $Z_N^{(\beta=2)}\{t\}$ is a τ -function of the Kadomtsev-Petviashvili (KP) hierarchy [2, 3, 4, 5]. In fact, the integral (1) can be identified as a continuum limit of the multisoliton solution of the hierarchy. This fact may be easily observed if we rewrite $Z_N^{(\beta=2)}$ in the following form:

$$Z_N^{(\beta=2)} = N! \det \left[\int_{-\infty}^{\infty} x^{j+k-2} e^{\eta(x,t)} dx \right]_{j,k=1,\dots,N}. \quad (2)$$

It may be also clear that the quantity $Z_N^{(\beta=2)}$ satisfies the bilinear Toda equation

$$Z_N^{(\beta=2)} \cdot \partial_{t_1}^2 Z_N^{(\beta=2)} - \left(\partial_{t_1} Z_N^{(\beta=2)} \right)^2 + Z_{N+1}^{(\beta=2)} \cdot Z_{N-1}^{(\beta=2)} = 0. \quad (3)$$

Here, we give other examples of such kinds of relationship. The orthogonal case $\beta = 1$ and the symplectic case $\beta = 4$ lead to τ -functions of the coupled KP hierarchy proposed

*Present Address: Department of Mathematical Sciences, Waseda University, Ohkubo 3-4-1, Shinjuku-ku, Tokyo 161-8555, Japan

by Hirota and Ohta [6]. The coupled KP hierarchy is an extension of the KP hierarchy. The first few bilinear equations of the hierarchy are

$$(D_1^4 - 4D_1D_3 + 3D_2^2)\tau \cdot \tau = 24\bar{\sigma}\sigma, \quad (4a)$$

$$(D_1^3D_2 + 2D_2D_3 - 3D_1D_4)\tau \cdot \tau = 12D_1\bar{\sigma} \cdot \sigma, \quad (4b)$$

$$(D_1^3 + 2D_3 + 3D_1D_2)\sigma \cdot \tau = 0, \quad (4c)$$

$$(D_1^4 - 4D_1D_3 - 3D_2^2 - 6D_4)\sigma \cdot \tau = 0, \quad (4d)$$

$$(D_1^3 + 2D_3 - 3D_1D_2)\bar{\sigma} \cdot \tau = 0, \quad (4e)$$

$$(D_1^4 - 4D_1D_3 - 3D_2^2 + 6D_4)\bar{\sigma} \cdot \tau = 0, \quad (4f)$$

where the Hirota bilinear operators are defined as

$$D_m^k D_n^l f \cdot g = \left(\frac{\partial}{\partial t_m} - \frac{\partial}{\partial t'_m} \right)^k \left(\frac{\partial}{\partial t_n} - \frac{\partial}{\partial t'_n} \right)^l f(t_1, t_2, \dots) g(t'_1, t'_2, \dots) \Big|_{t'=t}.$$

If we set

$$u = 2 \frac{\partial^2}{\partial t_1^2} \log \tau, \quad v = \sigma / \tau, \quad \bar{v} = \bar{\sigma} / \tau,$$

we obtain the coupled KP equation from the bilinear equations (4a), (4c), (4e) as

$$\frac{\partial}{\partial t_1} \left(4 \frac{\partial u}{\partial t_3} - 6u \frac{\partial u}{\partial t_1} - \frac{\partial^3 u}{\partial t_1^3} \right) - 3 \frac{\partial^2 u}{\partial t_2^2} + 24 \frac{\partial^2}{\partial t_1^2} (v\bar{v}) = 0, \quad (5a)$$

$$2 \frac{\partial v}{\partial t_3} + 3u \frac{\partial v}{\partial t_1} + \frac{\partial^3 v}{\partial t_1^3} + 3 \left(\frac{\partial^2 v}{\partial t_1 \partial t_2} + v \int^{t_1} \frac{\partial u}{\partial t_2} dt_1 \right) = 0, \quad (5b)$$

$$2 \frac{\partial \bar{v}}{\partial t_3} + 3u \frac{\partial \bar{v}}{\partial t_1} + \frac{\partial^3 \bar{v}}{\partial t_1^3} + 3 \left(\frac{\partial^2 \bar{v}}{\partial t_1 \partial t_2} + \bar{v} \int^{t_1} \frac{\partial u}{\partial t_2} dt_1 \right) = 0. \quad (5c)$$

We note that the coupled KP hierarchy reduces to the ordinary KP hierarchy by setting $\sigma = 0$ or $\bar{\sigma} = 0$. In that case, the coupled KP equations (3) reduce to the KP equation.

While the solutions of the KP hierarchy are expressed by determinants, the solutions of the coupled KP hierarchy are expressed in terms of pfaffians [6]. We denote a pfaffian associated with the $2N \times 2N$ antisymmetric matrix $A_{2N} = [a_{ij}]_{1 \leq i, j \leq 2N}$ as

$$\text{Pf}[A_{2N}] = \sum' \text{sgn} \begin{pmatrix} 1 & 2 & \cdots & 2n \\ j_1 & j_2 & \cdots & j_{2n} \end{pmatrix} a_{j_1 j_2} a_{j_3 j_4} \cdots a_{j_{2n-1} j_{2n}},$$

where the summation is taken over all $(j_1, \dots, j_{2n})$ such that $j_1 < j_3 < \cdots < j_{2n-1}$ and $j_1 < j_2, \dots, j_{2n-1} < j_{2n}$. Using this notation, solutions of the bilinear equations of the coupled KP hierarchy can be constructed as

$$\tau = \text{Pf}[A_{2N}], \quad \sigma = \text{Pf}[A_{2N-2}], \quad \bar{\sigma} = \text{Pf}[A_{2N+2}], \quad (6)$$

where the elements $a_{l,m}$ satisfy the differential rules

$$\frac{\partial}{\partial t_n} a_{l,m} = a_{l+n,m} + a_{l,m+n}, \quad (7)$$

for $n = 1, 2, \dots$. An example of the elements $a_{l,m}$ which satisfy eq. (7) is

$$a_{l,m} = \sum_k c_k (p_k^{l-1} q_k^{m-1} - p_k^{m-1} q_k^{l-1}) \exp [\eta(p_k, t) + \eta(q_k, t)]. \quad (8)$$

This τ -function corresponds to a multisoliton solution. If τ , σ and $\bar{\sigma}$ are of the form in eq. (6), the bilinear equations (2a-f) reduce to algebraic identities of pfaffians [6].

To obtain the relationship between the matrix integral and the coupled KP hierarchy, we use the equations attributable to de Bruijn [7]:

$$\begin{aligned} & \int \cdots \int_{x_1 \leq \cdots \leq x_N} \det [\phi_j(x_k)]_{j,k=1,\dots,N} dx_1 \cdots dx_N \\ &= \text{Pf} \left[\int \int \text{sgn}(y-x) \phi_j(x) \phi_k(y) dy dx \right]_{j,k=1,\dots,N} \\ &= \text{Pf} \left[\int \int_{x < y} (\phi_j(x) \phi_k(y) - \phi_j(y) \phi_k(x)) dy dx \right]_{j,k=1,\dots,N}, \end{aligned} \quad (9)$$

$$\begin{aligned} & \int \cdots \int \det [\phi_j(x_k), \psi_j(x_k)]_{j=1,\dots,2N, k=1,\dots,N} dx_1 \cdots dx_N \\ &= N! \text{Pf} \left[\int (\phi_j(x) \psi_k(x) - \phi_k(x) \psi_j(x)) dx \right]_{j,k=1,\dots,2N}. \end{aligned} \quad (10)$$

We note that Tracy and Widom utilized these formulas to obtain the Fredholm determinant expressions of similar matrix integrals [8]. Using these formulas, we first consider the orthogonal matrix model. Applying eq. (9) and the Vandermonde determinant

$$\prod_{j>k} (x_j - x_k) = \det [x_k^{j-1}]_{j,k=1,\dots,N},$$

we can rewrite $Z_N^{(\beta=1)}\{t\}$ in terms of a pfaffian as

$$\begin{aligned} Z_N^{(\beta=1)}\{t\} &= \int \cdots \int_{x_1 \leq \cdots \leq x_N} \prod_{1 \leq k < j \leq N} (x_j - x_k) \exp \left[\sum_{i=1}^N \eta(x_i, t) \right] dx_1 \cdots dx_N \\ &= \text{Pf} \left[\int \int_{x < y} (x^{j-1} y^{k-1} - y^{j-1} x^{k-1}) \exp [\eta(x, t) + \eta(y, t)] dx dy \right]_{j,k=1,\dots,N}. \end{aligned} \quad (11)$$

Because the elements of eq. (11) may be considered as a continuum limit of the multisoliton solution given by eq. (8), they satisfy the differential rules given by eq. (7). Hence, we conclude that $Z_N^{(\beta=1)}\{t\}$ is a τ -function of the coupled KP hierarchy, i.e., $Z_N^{(\beta=1)}\{t\}$ solves the bilinear equations (2a-f).

We then consider the symplectic case. Toward this aim, we prepare another formula. Consider the twofold Vandermonde determinant

$$\det [x_k^{j-1}, y_k^{j-1}]_{j=1,\dots,2N, k=1,\dots,N} = \prod_{j>k} (x_j - x_k) \cdot \prod_{j>k} (y_j - y_k) \cdot \prod_{j,k=1}^N (y_j - x_k). \quad (12)$$

Differentiating with respect to each y_k and setting $y_k = x_k$, we obtain the following equation [8]:

$$\prod_{j < k} (x_j - x_k)^4 = \det [x_k^j, (j-1)x_k^j]_{j=1,\dots,2N, k=1,\dots,N}. \quad (13)$$

Applying eqs. (10) and (13), we can rewrite $Z_N^{(\beta=4)}\{t\}$ in terms of a pfaffian as

$$\begin{aligned} Z_N^{(\beta=4)}\{t\} &= \frac{1}{N!} \int_{-\infty}^{\infty} \cdots \int_{-\infty}^{\infty} \prod_{1 \leq j < k \leq N} (x_j - x_k)^4 \exp \left[\sum_{i=1}^N \eta(x_i, t) \right] dx_1 \cdots dx_N \\ &= \text{Pf} \left[\int_{-\infty}^{\infty} (k-j) x^{k+j-3} \exp [2\eta(x, t)] dx \right]_{j,k=1, \dots, N}. \end{aligned} \quad (14)$$

Because the elements of eq. (14) satisfy eq. (7), $Z_N^{(\beta=4)}\{t\}$ is another example of a τ -function of the coupled KP hierarchy.

Lastly, we have demonstrated that the quantities

$$u^{(\beta)} = 2 \frac{\partial^2}{\partial t_1^2} \log Z_N^{(\beta)}\{t\}, \quad v^{(\beta)} = Z_{N-1}^{(\beta)}\{t\} / Z_N^{(\beta)}\{t\}, \quad \bar{v}^{(\beta)} = Z_{N+1}^{(\beta)}\{t\} / Z_N^{(\beta)}\{t\},$$

with $\beta = 1, 4$, satisfy the equations of the hierarchy such as the coupled KP equations (4). In the case $\beta = 2$, nonperturbative properties of the “specific heat” $u = 2\partial_{t_1}^2 \log Z_N^{(\beta=2)}\{t\}$ can be evaluated directly using the KP equation [9]. However, the properties of the coupled KP hierarchy are not as well understood as those of the ordinary KP hierarchy. For instance, Lax formulations and conserved quantities for the coupled hierarchy have not yet been obtained. Furthermore, although $Z_N^{(\beta=2)}$ satisfies the bilinear Toda equation (3), the corresponding facts for the cases $\beta = 1, 4$ are not known to date. We hope to report them in a forthcoming article.

Acknowledgments

The author gratefully acknowledges Dr. Yasuhiro Ohta for discussions and helpful comments. This work was supported by J.S.P.S. Research Fellowships for Young Scientists.

References

- [1] M. Kaku: *Strings, Conformal Fields, and Topology* (Springer, NY, 1991).
- [2] A. Morozov: Phys. Usp. **37** (1994) 1-55.
- [3] A. Marshakov, A. Mironov and A. Morozov: Phys. Lett. **B265** (1991) 99-107.
- [4] M. Mulase: In *Topology, Geometry and Field Theory* (World Sci. Publishing, River Edge, NJ, 1994) pp. 111-127.
- [5] M. Mulase: In *Perspectives in Mathematical Physics*, Conf. Proc. Lecture Notes Math. Phys. III (Internat. Press, Cambridge, MA, 1994) pp. 151-217.
- [6] R. Hirota and Y. Ohta: J. Phys. Soc. Jpn. **60** (1991) 798-809.
- [7] N. G. de Bruijn: J. Indian Math. Soc. **19** (1955) 133-151.
- [8] C. Tracy and H. Widom: *Correlation Functions, Cluster Functions and Spacing Distributions for Random Matrices*, preprint (solv-int/9804004).
- [9] V. A. Kazakov, I. K. Kostov and N. Nekrasov: *D-particles, Matrix Integrals and KP Hierarchy*, preprint (hep-th/9810035).